

Who Can Get a Third Dose of COVID-19 Vaccine?



This is an additional dose.

Studies indicate some immunocompromised people don't always build the same level of immunity after vaccination the way non-immunocompromised people do.

Only for those 12 and older who are immunocompromised.

12-17 year olds can only receive Pfizer.


Only for immunocompromised who have previously received Pfizer or Moderna.

No additional doses are approved for those that received Johnson and Johnson.

What to Know:



Must be at least 28 days since you received the second dose of Pfizer or Moderna.



You should receive the same vaccine you previously received.

(ie. If your first two doses were Pfizer your third would be Pfizer.)



If you do not meet the criteria below, you can NOT get an additional dose, at this time.

Who is Considered Immunocompromised

Including, but not limited to:

- Active treatment for **solid tumor** and **hematologic malignancies**
- Receipt of **solid-organ transplant** and taking immunosuppressive therapy
- Receipt of CAR-T-cell or hematopoietic **stem cell transplant** (within 2 years of transplantation or taking immunosuppression therapy)
- Moderate or severe primary **immunodeficiency** (e.g., DiGeorge syndrome, Wiskott-Aldrich syndrome)
- Advanced or untreated **HIV infection**

Active Treatment With:

- High-dose corticosteroids (i.e., ≥ 20 mg prednisone or equivalent per day),
- Alkylating agents
- Antimetabolites
- Transplant-related immunosuppressive drugs
- Cancer chemotherapeutic agents classified as severely immunosuppressive
- Tumor-necrosis (TNF) blockers
- Other biologic agents that are immunosuppressive or immunomodulatory